

REPORT DOCUMENTATION PAGE

FORM AD-266-1
OMB No. 0704-0188

AD-A266 250



JRT DATE

3 REPORT TYPE AND DATES COVERED

FINAL 01 Jan 89 TO 31 Dec 92

5 FUNDING NUMBERS

AFOSR-89-0245
61102F
2312
A2INVOLVEMENT OF LIPID METABOLISM IN CHEMICAL TRANSMISSION
PROCESSES AT MOSSY FIBER SYNAPSES

AUTHOR

Dr Robert V. Dorman

PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Kent State University
Dept of Biological Sciences
P.O. Box 5190
Kent, OH 44242-00018 PERFORMING ORGANIZATION
REPORT NUMBER

SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFOSR/NL
110 Duncan Avenue, Suite B115
Bolling AFB DC 20332-0001SPONSORING MONITORING
AGENCY REPORT NUMBER

Dr Haddad

SUPPLEMENTARY NOTES

12 DISTRIBUTION AVAILABILITY STATEMENT

Approved for public release;
distribution unlimited

93-14677



13 ABSTRACT (Maximum 200 words)

Isolated hippocampal mossy fiber synaptosomes were used to assess the relationships between membrane lipid metabolism and the evoked release of the excitatory amino acid neurotransmitter glutamate. A variety of metabolic parameters were investigated in order to develop a comprehensive model for the mechanisms of transmitter release. Mossy fiber terminals were radiolabeled with [3H] arachidonic acid and the effects of membrane depolarization and calcium influx on the labeling of the component lipid pools were determined. It was observed that depolarization and Ca²⁺ influx stimulated the accumulation of unesterified arachidonate. This effect was correlated with increased production of prostaglandins, in particular PGF_{2a} and could be blocked with lipase inhibitors and Ca²⁺ channel blockers. Prostaglandin production was also blocked by cyclooxygenase and lipoxxygenase inhibitors. We also observed that exogenous arachidonate stimulates the release of endogenous glutamate from mossy fiber terminals, as well as the mobilization of intraterminal free Ca²⁺. In addition, we identified two naturally occurring products which act synergistically to activate protein kinase C and facilitate the evoked release of glutamate.

14 SUBJECT TERMS

15. NUMBER OF PAGES

16 PRICE CODE

17 SECURITY CLASSIFICATION
OF REPORT

(U)

18 SECURITY CLASSIFICATION
OF THIS PAGE

(U)

19 SECURITY CLASSIFICATION
OF ABSTRACT

(U)

20 LIMITATION OF ABSTRACT

(UL)

FINAL TECHNICAL REPORT

TITLE: INVOLVEMENT OF LIPID METABOLISM IN CHEMICAL TRANSMISSION
PROCESSES AT MOSSY FIBER SYNAPSES

GRANT #: AFOSR-89-0245

PRINCIPAL INVESTIGATOR: Robert V. Dorman

DATE: MAY 24, 1993

SUMMARY

Isolated hippocampal mossy fiber synaptosomes were used to assess the relationships between membrane lipid metabolism and the evoked release of the excitatory amino acid neurotransmitter glutamate. A variety of metabolic parameters were investigated, in order to develop a comprehensive model for the mechanisms of transmitter release. Mossy fiber terminals were radiolabeled with [^3H]arachidonic acid and the effects of membrane depolarization and calcium influx on the labeling of the component lipid pools were determined. It was observed that depolarization and Ca^{2+} influx stimulated the accumulation of unesterified arachidonate. This effect was correlated with increased production of prostaglandins, in particular $\text{PGF}_{2\alpha}$ and could be blocked with lipase inhibitors and Ca^{2+} channel blockers. Prostaglandin production was also blocked by cyclooxygenase and lipoxygenase inhibitors. We also observed that exogenous arachidonate stimulates the release of endogenous glutamate from mossy fiber terminals, as well as the mobilization of intraterminal free Ca^{2+} . In addition, we identified two naturally occurring products which act synergistically to activate protein kinase C and facilitate the evoked release of glutamate.

RESEARCH OBJECTIVES

The general objective of this research was to examine the role that membrane lipids play in the mechanisms related to neurotransmitter release from a central synapse. Mossy fiber synaptosomes were isolated from rat hippocampus, because they play a role in learning and memory. In order to investigate the relationships between lipid metabolism and synaptic functions, some specific objectives were outlined.

Specific Objectives

1. Use [^3H]arachidonic acid to radiolabel nerve terminal lipids.
2. Determine the effects of membrane depolarization on arachidonate metabolism.
3. Assess the importance of external calcium on resting and depolarization-induced changes in arachidonate metabolism.
4. Investigate the possible mechanisms of arachidonate accumulation in response to depolarizing conditions.
5. Measure prostaglandin accumulation in response to membrane depolarization: assess effects of calcium antagonists and metabolic inhibitors.
6. Correlate the observed alterations in arachidonate and prostaglandin metabolism with the actual release of neurotransmitters and the accumulation of presynaptic calcium.

Accession No.	
NTIS	J
DTIC	
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	
61	
62	
63	
64	
65	
66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	
77	
78	
79	
80	
81	
82	
83	
84	
85	
86	
87	
88	
89	
90	
91	
92	
93	
94	
95	
96	
97	
98	
99	
100	

DTIC QUALITY INSPECTED 2, A-1

STATUS OF THE RESEARCH

The experimental work carried-out was designed to fit within the specific objectives listed above. A summary of the research follows.

1. Radiolabeling of glomerular lipids with [^3H]arachidonate:

Attempts were made to isotopically label hippocampal mossy fiber terminals in vitro. We found that arachidonate was incorporated into complex lipids and the uptake was linear with time. This method was then used for the studies of arachidonate metabolism.

2. Effects of membrane depolarization on arachidonate metabolism in the isolated mossy fiber synaptosomes:

Nerve terminals lipids were radiolabeled in vitro as described above. The labeled membranes were then used to assess the effects of membrane depolarization on the metabolic flux of arachidonate. We found that depolarization of the membranes with KCl, veratridine or 4-aminopyridine stimulated the accumulation of unesterified arachidonate. We suggest that the accumulation of unesterified arachidonate is an important response of nerve terminal membranes to depolarization and that such an effect is related to stimulus-secretion coupling.

3. Role of calcium in depolarization-induced arachidonate accumulation:

It is accepted dogma that the release of neurotransmitters requires the presence of external calcium. Therefore, we investigated the involvement of calcium in the evoked release of arachidonate. We found substantial inhibition of arachidonate accumulation in depolarized nerve terminals when Ca^{2+} was omitted from the incubation medium. We also found that verapamil, a Ca^{2+} channel blocker, inhibited the evoked release of arachidonate. The importance of external calcium for arachidonate accumulation was substantiated when the calcium ionophore A23187 was used to mimic the depolarization-induced effects on arachidonate metabolism when only the ionophore and external calcium were present. We interpret these results to show that the depolarization-induced accumulation of arachidonate is a calcium dependent process.

4. Prostaglandin production by hippocampal mossy fiber synaptosomes:

We previously found that there is an accumulation of arachidonate in stimulated terminals. So, we examined the relationship between arachidonate availability and the production of one of its major metabolites $\text{PGF}_{2\alpha}$. We observed that membrane depolarization enhanced PGF synthesis and this effect was blocked by calcium channel blockers and phospholipase inhibitors. In addition, we found that receptor activation with certain neurotransmitter agonists stimulated PG accumulation. Although we found no direct link between PG production and the secretion of glutamate from mossy fiber nerve endings, we have suggested that PG may modulate

transmitter release.

5. Correlations between observed alterations in lipid metabolism and the evoked release of neurotransmitters:

We were able to correlate the observed changes in membrane metabolism with the ultimate effect, the actual evoked release of neurotransmitters. This has provided us with the ability to make rapid advances in our understanding of release mechanisms. A summary of some of the correlative studies follow.

First, we showed that membrane depolarization evokes the release of glutamate from mossy fiber nerve terminals, as well as the accumulation of unesterified arachidonate and presynaptic calcium. Second, we found that the presence of exogenous arachidonate causes a dose-dependent release of neurotransmitter and this process is calcium-independent. The fact that the arachidonate-stimulated neurotransmitter release is calcium-independent has led us to suggest that the calcium requirement for transmitter release may be related to the accumulation of arachidonate. Third, we found that inhibition of the conversion of free arachidonate to prostaglandins with ibuprofen enhanced the depolarization- and arachidonate-evoked release of glutamate. Fourth, we observed that an oxidized derivative of arachidonate 12-HETE inhibited depolarization-induced glutamate release. We used the above information to propose that arachidonate provides an excitatory, presynaptic signal to stimulate glutamate release and that this effect is opposed by its 12-lipoxygenase products.

6. Mechanism of arachidonate modulation of evoked glutamate release.

We reported that exogenous arachidonate enhanced the amount of glutamate release from mossy fiber terminals induced by depolarization with KCl. More recently, we found that arachidonate and the diacylglycerol analog oleoyl-acetyl-glycerol acted synergistically to enhance the evoked release of glutamate. We found the same synergism with regards to the evoked accumulation of presynaptic calcium.

Both these effects were blocked by the protein kinase C inhibitor staurosporine. Therefore, we suggested that the two lipid metabolites work in concert to activate protein kinase C and potentiate the presynaptic responses to membrane depolarization.

PUBLICATIONS SUPPORTED BY AFOSR 89-0245:

Manuscripts:

1. Terrian, D.M., Damron, D.S., Dorman, R.V. and Gannon, R.L. (1989) Effects of calcium antagonists on the evoked release of dynorphin A 1-8 and availability of intraterminal calcium in rat hippocampal mossy fiber synaptosomes. *Neuroscience Letters* 106:322-327.
2. Freeman, E.J., Terrian, D.M. and Dorman, R.V. (1990) Presynaptic facilitation of glutamate release from isolated hippocampal mossy fiber nerve endings by arachidonic acid. *Neurochemical Research* 15:749-756.
3. Terrian, D.M., Dorman, R.V. and Gannon, R.L. (1990) Characterization of the presynaptic calcium channels involved in glutamate exocytosis from rat hippocampal mossy fiber synaptosomes. *Neuroscience Letters* 119:211-214.
4. Freeman, E.J., Damron, D.S., Terrian, D.M. and Dorman, R.V. (1991) 12-Lipoxygenase products attenuate the glutamate release and Ca^{2+} accumulation evoked by depolarization of hippocampal mossy fiber nerve endings. *Journal of Neurochemistry* 56:1079-1082.
5. Terrian, D.M., Dorman, R.V., Damron, D.S. and Gannon, R.L. (1991) Displacement of endogenous glutamate with D-aspartate: an effective strategy for reducing the calcium-independent component of glutamate release from synaptosomes. *Neurochemical Research* 16:35-41.
6. Dorman, R.V. (1991) PGF₂ synthesis in isolated cerebellar glomeruli: effects of membrane depolarization, calcium availability and phospholipase activity. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 42:233-240.
7. Dorman, R.V., Hamm, T.R.F., Damron, D.S. and Freeman, E.J. (1992) Modulation of glutamate release from hippocampal mossy fiber nerve endings by arachidonic acid and eicosanoids. In "Neurobiology of Essential Fatty Acids", ed. N.G. Bazan, Plenum Press, NY, pp. 121-136.
8. Separovic, D. and Dorman, R.V. (1993) Prostaglandin F₂alpha synthesis in the hippocampal mossy fiber synaptosomal preparation: I. Dependence on arachidonic acid, phospholipase A₂, calcium availability and membrane depolarization. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 48:127-137.
9. Damron, D.S. and Dorman, R.V. (1993) Involvement of phospholipase A₂ and arachidonic acid in the depolarization evoked accumulation of Ca^{2+} in hippocampal mossy fiber nerve endings. *Neurochemical Research* 18:1231-1237.
10. Separovic, D. and Dorman, R.V. (1993) Prostaglandin F₂alpha synthesis in the hippocampal mossy fiber synaptosomal preparation: II. Effects of receptor activation. *Prostaglandins, Leukotrienes and Essential Fatty Acids*. (submitted).

Abstracts:

1. Freeman, E.J., Terrian, D.M. and Dorman, R.V. (1989) Arachidonate evoked release of glutamate from mossy fiber terminals. Transactions of the American Society for Neurochemistry 20:267.
2. Terrian, D.M., Dorman, R.V. and Gannon, R.L. (1989) Effects of calcium antagonists on the evoked release of glutamate and dynorphin from hippocampal synaptosomes. Journal of Neurochemistry 52:S48.
3. Dorman, R.V., Freeman, E.J., Damron, D.S. and Terrian, D.M. (1989) Involvement of arachidonic acid in the release of endogenous glutamate from hippocampal mossy fiber synaptosomes. Conference on the Cell Biology of the Presynapse, Estoril, Portugal, p. 7.
4. Damron, D.S., Terrian, D.M. and Dorman, R.V. (1989) Calcium mobilization in hippocampal mossy fiber synaptosomes: phospholipase A₂ modulation. Neuroscience Abstracts 15:474.
5. Terrian, D., Damron, D., Gannon, R. and Dorman, R. (1989) Characterization of calcium channels in rat hippocampal mossy fiber synaptosomes. Neuroscience Abstracts 15:474.
6. Freeman, E.J., Terrian, D.M. and Dorman, R.V. (1990) Persistent enhancement of glutamate release by arachidonic acid. Transactions of the American Society for Neurochemistry 21:164.
7. Damron, D.S., Freeman, E.J., Terrian, D.M. and Dorman, R.V. (1990) Arachidonic acid-induced calcium mobilization in hippocampal mossy fiber synaptosomes. Neuroscience Abstracts 16:166.
8. Freeman, E.J., Damron, D.S., Terrian, D.M. and Dorman, R.V. (1990) Inhibition of glutamate release from hippocampal mossy fiber synaptosomes by 12-HETE. Neuroscience Abstracts 16:967.
9. Privette, T.H., Terrian, D.M., Zetts, D.A., Dorman, R.V. and Gannon, R.L. (1991) Kappa opioid autoregulation of the guinea pig hippocampal mossy fiber synapse. Transactions of the American Society for Neurochemistry 22:221.
10. Dorman, R.V., Damron, D.S., Freeman, E.J. and Terrian, D.M. (1991) Modulation of glutamate release from hippocampal mossy fiber nerve endings by arachidonic acid and eicosanoids. "Neurobiology of Essential Fatty Acids", Satellite to 13th International Society for Neurochemistry meeting, Cairns, Australia, July 10-12, 1991, p. 13.
11. Dorman, R.V., Agüero, E.G. and Bailey, A.S. (1992) Arachidonic acid and protein kinase C modulation of presynaptic calcium. Transactions of the American Society for Neurochemistry 23:194.
12. Separovic, D. and Dorman, R.V. (1992) Prostaglandin F_{2a} synthesis in hippocampal mossy fiber synaptosomes. Transactions of the American Society for Neurochemistry 23:149.
13. Dorman, R.V. and Zhang, L. (1993) Arachidonic acid and diglyceride facilitate evoked glutamate release. Transactions American Society for Neurochemistry 24:177.

PROFESSIONAL PERSONNEL

1. Nancy L. Edgehouse
July 1988 - 1992; Research Technician
2. Graduate Students receiving partial support through 89-0245:
Ernest J. Freeman, Ph.D., 1991
Derek S. Damron, Ph.D., 1991
Duska Separovic, Ph.D., 1992
Alex Bailey, MS, 1991